

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization

International Bureau

(43) International Publication Date
04 January 2018 (04.01.2018)



(10) International Publication Number
WO 2018/005588 A1

(51) International Patent Classification:

A01N 47/12 (2006.01) A01N 33/04 (2006.01)

(21) International Application Number:

PCT/US2017/039640

(22) International Filing Date:

28 June 2017 (28.06.2017)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/356,910 30 June 2016 (30.06.2016) US

(71) Applicant: **DOW GLOBAL TECHNOLOGIES LLC**
[US/US]; 2040 Dow Center, Midland, MI 48674 (US).

(72) Inventors: **FELDER, Patrick T.**; Bachtobelstrasse 3, CH-8810 Horgen (CH). **MURSET, Valerie**; Bachtobelstrasse 3, CH-8810 Horgen (CH). **VEZZOLI, Alessandro**; Bachtobelstrasse 3, CH-8810 Horgen (CH). **SZYMECZKO, Maciej**; Bachtobelstrasse 3, CH-8810 Horgen (CH). **YVON, Emmanuelle Christine**; Bachtobelstrasse 3, CH-8810 Horgen (CH).

(74) Agent: **EDWARDS, Tifani**; 400 Arcola Road, Collegeville, Pennsylvania 19426 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: SYNERGISTIC COMBINATION OF 3-iodo-2-propynyl-butylcarbamate AND DIAMINE

(57) Abstract: A synergistic antimicrobial composition containing 3-iodo-2-propynyl-butylcarbamate and diamine is provided. Also, a coating, a method of inhibiting the growth of or controlling the growth of microorganisms in an aqueous media, and a dry film containing 3-iodo-2-propynyl-butylcarbamate and diamine are further provided.



WO 2018/005588 A1

SYNERGISTIC COMBINATION OF
3-IODO-2-PROPYNYL-BUTYLCARBAMATE AND DIAMINE

This invention relates to combinations of biocides, the combinations having greater
5 activity than would be observed for the individual antimicrobial compounds.

Use of combinations of at least two antimicrobial compounds can broaden potential
markets, reduce use concentrations and costs, and reduce waste. In some cases, commercial
antimicrobial compounds cannot provide effective control of microorganisms, even at high
use concentrations, due to weak activity against certain types of microorganisms, or relatively
10 slow antimicrobial action, or instability under certain conditions such as high temperature and
high pH. Combinations of different antimicrobial compounds are sometimes used to provide
overall control of microorganisms or to provide the same level of microbial control at lower
use rates in a particular end use environment. Additionally, synergy has been found to be an
unpredictable phenomenon. Often like compounds display varying synergistic profiles when
15 combined with a particular active. It may be that no synergy is evidenced or it may be that
synergy exists but over a different synergistic range. Because of this observation, it is
difficult, if not impossible to draw conclusions regarding the synergistic profile of one
compound based upon the synergistic profile of a like compound. Thus more synergistic
combinations and their synergistic ranges must be discovered.

20 On such example of synergy is found in U.S. Pat. App. Pub. No. 2007/0078118. This
reference discloses synergistic combinations of N-methyl-1,2-benzisothiazolin-3-one (MBIT)
with other biocides. There still exists a need for additional combinations of antimicrobial
compounds having enhanced activity to provide effective control of microorganisms. The
problem addressed by this invention is to provide such combinations of antimicrobial
25 compounds.

In the present invention there is provided a synergistic antimicrobial composition
comprising 3-iodo-2-propynyl-butylcarbamate (also known as iodopropynyl butylcarbamate
and IPBC) (CAS registry number is 55406-53-6) and N-(3-Aminopropyl)-N-
dodecylpropane-1,3-diamine (also known as diamine) (CAS registry number is 2372-82-9).

30 The invention further provides a method of inhibiting the growth of or controlling the
growth of microorganisms in an aqueous media, the method comprising the step of adding a
synergistic antimicrobial composition comprising IPBC and diamine. Also, a coating, and a
dry film containing 3-iodo-2-propynyl-butylcarbamate and diamine are provided.

The following is a detailed description of the invention.

As used herein, the following terms have the designated definitions, unless the context clearly indicates otherwise.

5 The term "antimicrobial compound" refers to a compound capable of inhibiting the growth of or controlling the growth of microorganisms; antimicrobial compounds include bactericides, bacteriostats, fungicides, fungistats, algaecides and algistats, depending on the dose level applied, system conditions and the level of microbial control desired. Such term "antimicrobial compound" as used herein is synonymous with the term "biocide".

10 The term "microorganism" includes, for example, fungi (such as yeast and mold), bacteria and algae.

The following abbreviations are used throughout the specification: ppm = parts per million by weight (weight/weight), mL = milliliter, ATCC = American Type Culture Collection, SAG = Culture Collection of Algae at Goettingen University, CCAP = Culture Collection of Algae and Protozoa, DSMZ = Deutsche Sammlung von Mikroorganismen und Zellkulturen, and MIC = minimum inhibitory concentration.

20 Unless otherwise specified, temperatures are in degrees centigrade (°C), and references to percentages are by weight (wt.%). Percentages of antimicrobial compounds in the composition of this invention are based on the total weight of active ingredients in the composition, i.e., the antimicrobial compounds themselves, exclusive of any amounts of solvents, carriers, dispersants, stabilizers or other materials which may be present.

As used herein, "IPBC" is 3-Iodo-2-propynyl-butylcarbamate (CAS registry number 55406-53-6).

As used herein, "Diamine" is N-(3-Aminopropyl)-N-dodecylpropane-1,3-diamine (CAS registry number is 2372-82-9)

25 When a ratio is the herein to be "X:1 or higher," it is meant that the ratio is Y:1, where Y is X or greater, and when a ratio is the herein to be "X:1 or lower," it is meant that the ratio is Z:1, where Z is X or less. The same logic follows for ratios that are "1:X or higher" and "1:X or lower".

30 The present invention is a composition that contains both IPBC and diamine. It has been surprisingly found that compositions that contain both IPBC and diamine are synergistically effective as biocides. In the present invention, the weight ratio of the IPBC to diamine is from 1:10 to 10:1 alternatively from 1:1 to 1:10 and further alternatively from 10:1 to 1:1.

In some embodiments of the invention, the antimicrobial combination of this invention is useful for inhibiting the growth of or controlling the growth of microorganisms in an aqueous media. Such aqueous media includes but is not limited to industrial water and water containing/contaminated media, such as cooling water, air washer, heat exchangers, boiler water, pulp and paper mill water, other industrial process water media such as: ballast water, wastewater, metalworking fluids, oil and gas, latex, paint, coatings, adhesives, inks, tape joint compounds, pigment, water-based slurries, personal care and household products such as detergent, filtration systems (including reverse osmosis and ultrafiltration systems), toilet bowl, textiles, leather and leather production system, or a system used therewith. In one embodiment the antimicrobial composition is used as an in-can preservative.

Typically, the amount of the biocide combinations of the present invention for inhibiting the growth of or controlling the growth microorganisms is from 10 ppm to 5,000 ppm active ingredient. In some embodiments of the invention, the active ingredients of the composition are present in an amount of at least 20 ppm, alternatively at least 50 ppm, alternatively at least 100 ppm, alternatively at least 150 ppm, alternatively at least 200 ppm. In some embodiments, the active ingredients of the composition are present in an amount of no more than 2,000 ppm, alternatively no more than 1,000 ppm, alternatively no more than 500 ppm, alternatively no more than 400 ppm, alternatively no more than 300 ppm, alternatively no more than 250 ppm, alternatively no more than 200 ppm, alternatively no more than 100 ppm, alternatively no more than 50 ppm. Concentrations mentioned above are in a liquid composition containing the biocide combinations.

The present invention also encompasses a method for inhibiting the growth of or controlling the growth of microorganisms in the use areas described above, especially in in-can preservative applications, by incorporating the claimed biocide combination into the materials.

The composition of the present invention contains IPBC and diamine. It is contemplated that some embodiments may contain one or more additional antimicrobial compound.

The following are examples of the present invention.

The synergism of the biocides combination of the present invention was determined using the method described by Kull, F.C., et. al in *Applied Microbiology* 9:538-541 (1961). The formula to calculate the synergy index (SI) is

$$Qa/QA + Qb/QB = SI$$

Where

QA= concentration of compound A in ppm, acting alone produced an end point (growth / no growth)

5 Qa = concentration of compound A in ppm, in the mixture, which produced an end point (growth / no growth)

QB= concentration of compound B in ppm, acting alone produced an end point (growth / no growth)

Qb= concentration of compound B in ppm, in the mixture, which produced an end point (growth / no growth)

10 Synergism within two biocides is demonstrated when the SI has a value less than 1. The mixtures showed an additive effect if SI is equal to 1 and antagonistic if SI is greater than 1.

15 The Minimum Inhibitory Test (MIC) is designed to evaluate the lowest concentration of a biocide, biocide blend or biocide combination to prevent microorganisms from growing in a defined broth.

The MIC and synergy testing was carried out as follows:

1. The test was executed with a Hamilton MLStarPlus robot using automated turbidity reading with BioTek Synergy H4 plate reader.
- 20 2. Biocide plates were prepared in 2.2 ml deep well plates by transferring and diluting biocides from stock solutions to first row of the plates. The concentrations of biocides in stock bottles were adjusted to be 20x more concentrated than the highest desired concentration.
3. Then 15 subsequent serial dilutions with dilution factor 1.3 were performed resulting in 16 different concentrations for each system.
- 25 4. In the next step serially diluted biocide systems were transferred to the media blocks containing 850 μ l of tryptic soy broth, comprising, Casein (pancreatic digest) 17 g/L, Soya peptone (papaic digest) 3 g/L, Sodium chloride 5 g/L, Dipotassium phosphate 2.5 g/L, Dextrose 2.5 g/L ("TSB") containing different concentrations of Mowiol 18-88 partially hydrolyzed polyvinylacetate surfactant, commercially available from
30 Kuraray Europe GmbH in each well. In case of single biocide systems 100 μ l were transferred and in case of biocide combinations 50 μ l of each biocide dilution were transferred to the media, resulting in 950 μ l of final volume of media + biocides and

9.5 times dilution of the biocides from the biocide plate. At this point, the concentrations of all biocides in media were 1.053 x final concentration.

5. After preparation and mixing of the described systems, 3 aliquots of 190 μ l were prepared in 96-well microtiter plates.

5 6. Preparation of the microbe suspension:

Bacterial cultures:

Pseudomonas aeruginosa DSM # 939 ATCC# 15442

Staphylococcus aureus DSM # 799 ATCC# 6538

10 The culture was maintained as a glycerol stock at -80°C in cryovials. A cryovial was thawed and then 100 μ l spread on a TSB agar plate. After incubation for 1 day at 30°C the bacteria were harvested with buffer at pH 7.3. A total viable count on TSB plate was carried out and bacterial suspension was diluted in buffer in order to deliver ~2x10⁷ CFU/ml.

Yeast culture:

15 *Candida albicans* DSM #1386 ATCC# 10231

The cultures were maintained as glycerol stocks at -80°C in cryovials, are thawed and then 100 μ l spread on MEA (malt extract agar) petri dishes.

The yeast strain plates were incubated at 28°C for 1-2 days then harvested with buffer pH 5.0.

20 Based on total viable count results, the inoculum was prepared.

7. Each test sample (190 μ l) was inoculated with the 10 μ l of microbe suspension to provide a level of ~1 x 10⁶ CFU/ml of the yeast species.

8. The test samples were mixed and incubated at 30°C for 2 days (48 hours) when tested against bacteria and 3 days (72 hours), respectively, when tested against yeast.

9. Growth of the micro-organisms leads to turbidity after incubation, clarity indicates no growth. Reading of the results was carried out by measuring absorbance at 600 nm for each sample at the beginning of the test (t_{zero}) and after incubation ($t_{endpoint}$). $t_{endpoint}$ was chosen at 48 hours for bacteria and 72 hours for yeast. The difference in absorbance between $t_{endpoint}$ and t_{zero} was used to assign a score (“1” if $\Delta > 0.2$, confirming growth, and “0” if $\Delta \leq 0.2$, confirming no growth) from which the MIC values were derived. The lowest concentration that showed no growth (score of “0”) in the broth after incubation is taken as the MIC value.

The results of single biocide and combination of two biocides tested against bacteria and yeast are presented in Tables 1 and 2.

Table 1: MIC results for single biocides and combinations of two biocides (in ppm):

Active ingredients and ratio	<i>Staphylococcus aureus</i> (DSMZ#799)	<i>Pseudomonas aeruginosa</i> (DSMZ#939)	<i>Candida albicans</i> (DSMZ#1386)
IPBC (single)	21.6	1400.0	6.6
Diamine (single)	16.7	47.8	61.5
IPBC/Diamine 1:1	5.8 : 5.8	28.3 : 28.3	2.6 : 2.6
IPBC/Diamine 1:10	1.3 : 12.9	6.2 : 62.2	1.0 : 9.8
IPBC/Diamine 10:1	9.8 : 1.0	223.1 : 22.3	5.1 : 0.5

Table 2: Calculated synergy indices for the Combinations in Table 1

Active ingredients and ratio	<i>Staphylococcus aureus</i> (DSMZ#799)	<i>Pseudomonas aeruginosa</i> (DSMZ#939)	<i>Candida albicans</i> (DSMZ#1386)
IPBC/Diamine 1:1	0.62	0.61	0.44
IPBC/Diamine 1:10	0.83	1.30	0.31
IPBC/Diamine 10:1	0.51	0.63	0.78

CLAIMS

1. A synergistic antimicrobial composition comprising 3-iodo-2-propynyl-butylcarbamate and diamine.
2. The synergistic antimicrobial composition of claim 1, wherein the weight ratio of the 3-iodo-2-propynyl-butylcarbamate to diamine is from 1:10 to 10:1.
3. A method of inhibiting the growth of or controlling the growth of microorganisms in an aqueous media.
4. The method of claim 3 wherein the aqueous media is a coating.
5. A coating composition comprising the synergistic antimicrobial composition of claim 1.
6. A coating composition comprising the synergistic antimicrobial composition of claim 2.
7. A dry film made by a process comprising applying a layer of the coating composition of claim 4 to a substrate and drying the coating composition or allowing the coating composition to dry.

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2017/039640

A. CLASSIFICATION OF SUBJECT MATTER
INV. A01N47/12 A01N33/04
ADD.
According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED
Minimum documentation searched (classification system followed by classification symbols)
A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
EPO-Internal, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 03/062149 A2 (LONZA AG [US]; LONZA AG [CH]; LUDENSKY MICHAEL [US]; HILL CHRISTOPHER) 31 July 2003 (2003-07-31) example 1 claims 16-21	1-7
A,P	WO 2016/137830 A1 (DOW GLOBAL TECHNOLOGIES LLC [US]) 1 September 2016 (2016-09-01) the whole document	1-7

Further documents are listed in the continuation of Box C.

See patent family annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier application or patent but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search 4 August 2017	Date of mailing of the international search report 14/08/2017
---	---

Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Götz, Gerhard
--	--

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2017/039640

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 03062149	A2	31-07-2003	
		AU 2003237529 A1	02-09-2003
		US 2003228373 A1	11-12-2003
		WO 03062149 A2	31-07-2003

WO 2016137830	A1	01-09-2016	NONE
