

FIGURE 1
Scheme 1




FIGURE 2

Table 1
 Structures of (E)-stilbenes,^a

Product	R1	R2	Reference ^b	Product	R1	R2	Reference ^b
1	4-OCH ₃	2-F	21	29*	2-F	2,5-diF	-
2	-	2-F	21	30*	3-F	3,4-diF	-
3	-	3-F	21	31	2-F	2,6-diF	35
4	-	4-F	21	32	3-F	3-F	31
5	4-OCH ₃	3-F	21	33	3-F	4-F	36
6	4-OCH ₃	4-F	21	34*	3-F	2,5-diF	-
7	4-CH ₃	4-F	28	35*	3-F	2,4-diF	-
8	4-CH ₃	4-F	28	36*	2-F	2,4-diF	-
9	4-CH ₃	3-F	28	37*	4-CH ₃	2,4-diF	-
10*	2-F	2-OCH ₃	-	38	4-F	2,4-diF	39
11	3-OCH ₃	2-F	29	39	4-F	3,4-diF	39
12	3-OCH ₃	4-F	30	40*	4-F	2,5-diF	-
13	3-OCH ₃	3-F	31	41	-	2,3-diOCH ₃	21
14	3-F	3,4-diOCH ₃	32	42	-	2-OCH ₃	21
15	4-F	3,4-diOCH ₃	33	43	4-OCH ₃	2,4-diOCH ₃	21
16*	3-F	2,3-diOCH ₃	-	44	4-OCH ₃	2-Cl	21
17*	4-F	2,3-diOCH ₃	-	45	4-OCH ₃	3-OH	34
18*	4-F	2,5-diOCH ₃	-	46	4-OCH ₃	3-OCH ₃ , 4-OH	40
19	4-Cl	2-F	29	47	4-CH ₃	3-CF ₃	31
20	4-OH	2-F	34	48*	2,3-OCH ₃	4-CH ₃	-
21	-	2,6-diF	35	49*	3-OCH ₃	2-OCH ₃	-
22*	4-CH ₃	3,4-diF	-	50*	3-OCH ₃	2,5-diOCH ₃	-
23*	4-OCH ₃	2,5-diF	-	51	4-OCH ₃	3-NO ₂	41
24*	2-F	3,4-diF	-	52	-	3-NO ₂	42
25	2-F	3-F	36	53	4-CH ₃	3-NO ₂	43
26*	4-OCH ₃	3,4-diF	-	54*	2,3-diOCH ₃	4-CF ₃	-
27	2-F	2-F	37	55*	2,3-diOCH ₃	4-CH(CH ₃) ₂	-
28	4-F	2-F	38	56	2,4-diOCH ₃	2,5-diOCH ₃	44

^a Data for starred compounds is reported in the experimental section.

^b Melting points and NMR data of known compounds are in agreement with the literature data.

FIGURE 3

Table 2. Activation of Nrf2 by mono-fluoro *trans* stilbenes













Number	Structure	EC ₅₀ (μM)	Fold activation
Sulforaphane		1.2 ± 0.08	12.4
Resveratrol		5.4 ± 0.5	3.2
1 (LD55)		5.4 ± 0.3	14.9
2		0.7 ± 0.1	10.7
3		3.1 ± 0.4	4.9
4		12.4 ± 0.5	4.8
5		8.9 ± 0.9	3.6
6		4.2 ± 0.4	1.7
7		1.3 ± 0.2	1.1
8		0.8 ± 0.08	4.7
9		1.4 ± 0.2	10.1
10		3.1 ± 0.2	18.4

FIGURE 3 (Cont'd)











Number	Structure	EC ₅₀ (μM)	Fold activation
11		3.8 ± 0.2	14.4
12		6.0 ± 0.5	4.4
13		2.3 ± 0.5	15.1
14		9.5 ± 0.7	5.9
15		2.6 ± 0.5	1.6
16		1.9 ± 0.1	19.4
17		2.9 ± 0.1	16.9
18		5.1 ± 0.5	10.7
19		1.6 ± 0.2	5.3
20		11.4 ± 1.6	13.4

FIGURE 4

Table 3. Activation of Nrf2 by polyfluoro *trans* stilbenes












Number	Structure	EC ₅₀ (μM)	Fold activation
21		6.9 ± 0.1	11.4 ± 0.4
22		2.6 ± 0.18	10.2 ± 0.2
23		3.2 ± 0.5	17.0 ± 0.9
24		>15	11.5 ± 2.0 ^a
25		>15	8.5 ± 0.2 ^a
26		>15	4.6 ± 0.6 ^a
27		4.3 ± 0.4	13.9 ± 0.9
28		3.6 ± 0.3	12.7 ± 0.4
29		5.9 ± 0.2	22.3 ± 0.5
30		5.0 ± 0.1	12.2 ± 0.8
31		1.4 ± 0.5	21.2 ± 2.0

FIGURE 4 (Cont'd)

Table 3 (Cont'd)










Number	Structure	EC ₅₀ (μM)	Fold activation
32		1.0 ± 0.12	7.0 ± 0.7
33		0.65 ± 0.12	4.5 ± 0.4
34		0.3 ± 0.02	10.7 ± 1.0
35		0.45 ± 0.01	8.2 ± 0.2
36		9.6 ± 0.14	5.0 ± 0.8
37		2.0 ± 0.07	13.5 ± 0.2
38		0.9 ± 0.08	16.8 ± 1.8
39		>15	7.0 ± 0.5 ^a
40		2.9 ± 0.59	11.2 ± 0.1

FIGURE 5

Table 4. Activation of Nrf2 by non-fluoro *trans* stilbenes










Number	Structure	EC ₅₀ (μ M)	Fold activation
41		4.7 \pm 0.7	25.4
42		4.4 \pm 0.1	23.1
43		4.6 \pm 0.4	15.6
44		3.1 \pm 0.2	17.8
45		>15	28.1
46		3.7 \pm 0.5	27.7
47		1.5 \pm 0.5	19.2
48		1.1 \pm 0.2	16.1
49		4.4 \pm 0.2	29.2

FIGURE 5 (Cont'd)

Table 5 (Cont'd)








50		>15	15.7
51		3.3 ± 0.4	69.0
52		3.0 ± 0.4	18.6
53		3.7 ± 0.2	11.9
54		2.2 ± 0.1	65.5
55		2.7 ± 0.3	33.9
56		0.8 ± 0.03	12.9

FIGURE 6

EC₅₀ values for inhibition of NF-κB and activation of Nrf2 by resveratrol analogues (trans stilbenes) and by curcumin analogues (dienones)














Compound	Structure	EC ₅₀ (μM) NFκB	EC ₅₀ (μM)Nrf2
Resveratrol		20 ± 3	5.4±0.5
Curcumin		8.2±0.4	21±0.02
1 (LD55)		0.15 ± 0.1	5.4±0.3
5		1.1±0.6	8.9±0.9
6		3-5	4.2±0.4
7		3-5	1.3±0.2
8		0.77±0.3	0.8±0.08

FIGURE 6 (Cont'd)

Table 6 Continued

Compound	Structure	EC ₅₀ (μM) NFκB	EC ₅₀ (μM) Nrf2
9		0.63±0.5	1.4±0.2
11		3 - 5	3.8±0.2
12		3 - 5	6.0±0.5
44		1.5±0.03	3.1±0.2
dienone analogue of curcumin		4.1±0.3	3.3±0.5
Dienone analogue		9.6±0.7	9.9±0.6